

Abelianization of p-groups with derived subgroup of prime order

Guhan Venkat

III Year, Int. MS, Indian Institute of Science Education and Research, Pune.
guhan.venkat@gmail.com

Subject Class : GR

Mathematics Subject Classification : 20E34 (General Structure Theorems)

Abstract

In this paper, we arrive at the structure for $G^{ab}(= G/G')$ from a presentation for G . We also prove that the frattini subgroup of the abelianization is G^p or the quotient of G^p by the derived subgroup.

Key words : Abelianization, Frattini Subgroup, Class 2.

Introduction

Blackburn, S.R classified all p-groups, G , with derived subgroup of prime order in [1]. In this paper, we find the abelianization of such groups. We interchangably use G^{ab} and G/G' .

Let G be a p-group of order p^n where p is an odd prime and $n > 1$. Since G is nilpotent, $[G', G]$ is strictly contained in $G' \cong \mathbb{Z}_p$ and thus $[G', G] = 1$. Thus G has nilpotency class 2 $\Rightarrow G' \subseteq Z(G)$.

Basic Lemmas

Lemma 1

$$[x^p, y] = [x, y]^p = 1.$$

Proof: The first equality follows from "class 2" and the second from the fact $|G'| = p$.

Lemma 2

$$(xy)^p = x^p y^p [y, x]^{1/2(p)(p-1)} = x^p y^p.$$

Proof: The first equality is infact true for all integers and can be proved using induction. The second equality follows because p is an odd prime and 2 divides $(p-1)$ and from Lemma 1.

Structure of G^{ab}

Given $G = \langle x_1, x_2, \dots, x_k \rangle$, any element of $a \in G$ can be written as

$$a = \prod_{i=1}^k x_i^{e_i} \Rightarrow a^p = \prod_{i=1}^k (x_i^{e_i})^p = (x_i^p)^{e_i}$$

(from Lemma 2). Also, we have that $(xy)^p \in Z(G) \forall x, y \in G$.

Thus we have shown that if $G = \langle x_1, x_2, \dots, x_k \rangle$, then $G^p = \langle x_1^p, x_2^p, \dots, x_k^p \rangle$ and also that $G^p \subseteq Z(G)$. In particular G^p is a finite abelian group and hence all relations in G^p are power relations on the generators (from the Structure Theorem) which carry over from power relations in G .

From [2], it follows that

$$\Phi(G/N) \cong \Phi(G)/N$$

where $N \text{ char } G$ and $N \trianglelefteq \Phi(G)$

Taking $N = G' \cong \mathbb{Z}_p$, we have

$$\Phi(G/G') \cong \Phi(G)/G'$$

$$\Phi(G) = G^p G'$$

$$\Rightarrow \Phi(G/G') = G^p G' / G' \cong G^p / G^p \cap G' \text{ (from Second Isomorphism Theorem).}$$

In particular, when $\Phi(G) = G' \cong \mathbb{Z}_p$, then G/G' is elementary abelian. Such groups are known as generalized extraspecial and have been classified in [3].

Since $G^p \cap G' \trianglelefteq G'$, $G^p \cap G' \cong G'$ or is trivial.

Case (i)

$$G^p \cap G' = G' \Rightarrow \Phi(G/G') \cong G^p/G'.$$

Writting G^p and G' additively, one has the following presentation for G^p/G'

$$\begin{pmatrix} p^{\lambda_1} & & & \\ & p^{\lambda_2} & & \\ & & \ddots & \\ & & & p^{\lambda_k} \\ p & & \dots & \end{pmatrix} \begin{pmatrix} px_1 \\ px_2 \\ \vdots \\ px_k \end{pmatrix} = 0$$

where each x_i has order p^{λ_i+1} in G $0 \leq \lambda_i < n$

Note : We re-label the generators in a way that $G' = \langle [x_i, x_j] \rangle = \langle x_1^p \rangle$. (which can always be done from a power commutator presentation for G)

Computing the Smith Normal form for the matrix above, we get the structure of $\Phi(G^{ab})$ from which we get the structure of G^{ab} .

Case (ii)

$$G^p \cap G' = e \Rightarrow \Phi(G/G') \cong G^p$$

Accordingly we get the structure of the abelianization of G .

Given a presentation for G , one can easily find out if $G^p \cap G' = e$ from the ‘‘Commutator Relation’’ of G .

References

- [1] Blackburn, Simon. R, Groups of prime power order with derived subgroup of prime order, *J. Algebra* 219 (2) (1999) 625-657.
- [2] Hobby, Charles, The Frattini Subgroup of a p-group, *Pacific J. Math* 10 (1) (1960) 209-212.
- [3] Stancu, Radu, Almost all Generalized Extraspecial p-groups are Resistant, *J. Algebra* 249 (1) (2002) 120-126.